

NOTE: For problems involving motion in free fall, use $g = 9.80 \text{ m/s}^2$ unless otherwise specified.

Q1) A car travels 40 kilometers at an average speed of 80 km/h and then travels 40 kilometers at an average speed of 40 km/h. The average speed of the car for this 80 km trip is:

- A) 40 km/h
- B) 45 km/h
- C) 53 km/h**
- D) 60 km/h
- E) 80 km/h


Q2) A car starts from rest and goes down a slope with a constant acceleration of 5 m/s^2 . After 5 seconds the car reaches the bottom of the hill. What is its speed at the bottom of the hill?

- A) 1 m/s
- B) 12.5 m/s
- C) 25 m/s**
- D) 50 m/s
- E) 160 m/s


Q3) A baseball is thrown vertically into the air. The acceleration of the ball at its highest point is:

- A) zero
- B) g , down**
- C) g , up
- D) $2g$, down
- E) $2g$, up


Q4) The vectors \vec{a} , \vec{b} , and \vec{c} are related by $\vec{c} = \vec{a} - \vec{b}$. Which diagram below illustrates this relationship?




I



II




III



IV

- A) I.
- B) II.
- C) III.**
- D) IV.
- E) None of these

Q5) In the diagram, \vec{A} has magnitude 12 m and \vec{B} has magnitude 8 m. The x component of $\vec{A} + \vec{B}$ is approximately:




- A) 1.5 m
- B) 4.5 m
- C) 12 m**
- D) 15 m
- E) 20 m

Q6) A constant force of 8.0 N is exerted for 4.0 s on a 16-kg object initially at rest. The change in speed of this object will be:

- A) 0.5 m/s
- B) 2 m/s**
- C) 4 m/s
- D) 8 m/s
- E) 32 m/s

Q7) Two blocks weighing 250 N and 350 N respectively, are connected by a string that passes over a massless pulley as shown. The tension in the string is:



- A) 210 N
- B) 290 N**
- C) 410 N
- D) 500 N
- E) 4900 N

Q8) A 32-N force, parallel to the incline, is required to push a certain block at constant velocity up a frictionless incline that is 30° above the horizontal. The mass of the block is:

- A) 3.3 kg
- B) 3.8 kg
- C) 5.7 kg
- D) 6.5 kg**
- E) 160 kg

Q9) A 12-kg block rests on a horizontal surface and a boy pulls on it with a force that is 30° below the horizontal. If the coefficient of static friction is 0.40, the minimum magnitude force he needs to start the block moving is:

- A) 44 N
- B) 47 N
- C) 54 N
- D) 56 N
- E) 71 N**

Q10) A 5.0-kg block is resting on a horizontal plank. The coefficient of static friction is 0.50 and the coefficient of kinetic friction is 0.40. After one end of the plank is raised so the plank makes an angle of 25° with the horizontal, the force of friction is approximately:

- A) 0 N
- B) 20 N
- C) 21 N**
- D) 22 N
- E) 44 N

Q11) A 5.0-kg block is resting on a horizontal plank. The coefficient of static friction is 0.50 and the coefficient of kinetic friction is 0.40. After one end of the plank is raised so the plank makes an angle of 30° with the horizontal, the force of friction is:

- A) 0 N
- B) 17 N**
- C) 20 N
- D) 25 N
- E) 49 N

Q12) A 5.0-kg block is on an incline that makes an angle of 30° with the horizontal. If the coefficient of static friction is 0.50, the minimum force that can be applied parallel to the plane to hold the block at rest is:

- A) 0 N
- B) 3.3 N**
- C) 21 N
- D) 25 N
- E) 46 N


Q13) A 5.0-kg block is on an incline that makes an angle 30° with the horizontal. If the coefficient of static friction is 0.5, the maximum force that can be applied parallel to the plane without moving the block is:

- A) 0 N
- B) 3.3 N
- C) 21 N
- D) 46 N**
- E) 55 N

Q14) Block A, with mass m_A , is initially at rest on a horizontal floor. Block B, with mass m_B , is initially at rest on the horizontal top surface of A. The coefficient of static friction between the two blocks is μ_s . Block A is pulled with a horizontal force. It begins to slide out from under B if the force is greater than:

- A) $m_A g$
- B) $m_B g$
- C) $\mu_s m_A g$
- D) $\mu_s m_B g$
- E) $\mu_s (m_A + m_B)g$**

Q15) The system shown remains at rest. The force of friction on the block on the slope is:



- A) 4 N
- B) 8 N**
- C) 12 N
- D) 16 N
- E) 20 N

Q16) A 1000-kg airplane moves in straight flight at constant speed. The force of air friction is 1800 N. The net force on the plane is:

- A) 0 N**
- B) 11600 N
- C) 1800 N
- D) 9800 N
- E) none of these

Q17) A rock is dropped from the top of a vertical cliff and takes 3.00 s to reach the ground below the cliff. A second rock is thrown vertically from the cliff, and it takes this rock 2.00 s to reach the ground below the cliff from the time it is released. With what velocity was the second rock thrown, assuming no air resistance?

- A) 4.76 m/s upward
- B) 5.51 m/s downward
- C) 12.3 m/s upward
- D) 4.76 m/s downward
- E) 12.3 m/s downward**


Q18) A test rocket is fired straight up from rest with a net acceleration of 20.0 m/s^2 . After 4.00 seconds the motor turns off, but the rocket continues to coast upward with no appreciable air resistance. What maximum elevation does the rocket reach?

- A) 487 m
- B) 327 m
- C) 320 m
- D) 408 m
- E) 160 m

Q19) A ball is projected upward at time $t = 0.00 \text{ s}$, from a point on a roof 70 m above the ground and experiences negligible air resistance. The ball rises, then falls and strikes the ground. The initial velocity of the ball is 28.5 m/s . Consider all quantities as positive in the upward direction. The velocity of the ball when it is 39 m above the ground is closest to


- A) -38 m/s .
- B) -30 m/s .
- C) -23 m/s .
- D) -15 m/s .
- E) -45 m/s .

Q20) A 4.00-kg block rests between the floor and a 3.00-kg block as shown in the figure. The 3.00-kg block is tied to a wall by a horizontal rope. If the coefficient of static friction is 0.800 between each pair of surfaces in contact, what horizontal force F must be applied to the 4.00-kg block to make it move?




- A) 16.2 N
- B) 54.9 N
- C) 21.1 N
- D) 23.5 N
- E) 78.4 N

Q21) A rope pulls on the lower block in the figure with a tension force of 20 N. The coefficient of kinetic friction between the lower block and the surface is 0.16. The coefficient of kinetic friction between the lower block and the upper block is also 0.16. The pulley has no appreciable mass or friction. What is the acceleration of the 2.0 kg block?




- A) 4.1 m/s²
- B) 5.1 m/s²
- C) 8.4 m/s²
- D) 9.2 m/s²

Q22) A system comprised blocks, a light frictionless pulley, and connecting ropes is shown in the figure. The 9.0-kg block is on a perfectly smooth horizontal table. The surfaces of the 12-kg block are rough, with $\mu_k = 0.30$ between the two blocks. If the 5.0-kg block accelerates downward when it is released, find its acceleration.




- A) 1.0 m/s²
- B) 1.2 m/s²
- C) 1.4 m/s²
- D) 1.6 m/s²
- E) 1.8 m/s²

Q23) Block A of mass 5.0 kg and block X are attached to a rope which passes over a pulley, as shown in the figure. An 80-N force P is applied horizontally to block A, keeping it in contact with a rough vertical face. The coefficients of static and kinetic friction between the wall and block A are $\mu_s = 0.40$ and $\mu_k = 0.30$. The pulley is light and frictionless. The mass of block X is adjusted until block A moves upward with an acceleration of 1.6 m/s^2 . What is the mass of block X?




- A) 9.9 kg
- B) 9.3 kg
- C) 8.7 kg
- D) 8.1 kg
- E) 7.5 kg

Q24) Block A of mass 8.0 kg and block X are attached to a rope that passes over a pulley. A 50-N force P is applied horizontally to block A, keeping it in contact with a rough vertical face. The coefficients of static and kinetic friction between the wall and block A are $\mu_s = 0.40$ and $\mu_k = 0.30$. The pulley is light and frictionless. In the figure, the mass of block X is adjusted until block A descends at constant velocity of 4.75 cm/s when it is set into motion. What is the mass of block X?



- A) 6.5 kg
- B) 7.2 kg
- C) 8.0 kg
- D) 8.8 kg
- E) 9.5 kg

Q25) Three objects are connected as shown in the figure. The strings and frictionless pulleys have negligible masses, and the coefficient of kinetic friction between the 2.0-kg block and the table is 0.25. What is the acceleration of the 2.0-kg block?



- A) 2.5 m/s^2
- B) 1.7 m/s^2
- C) 3.2 m/s^2
- D) 4.0 m/s^2