

1. Starting with a sample of pure ^{66}Cu , $7/8$ of it decays into Zn in 15 minutes. The corresponding half-life is:

- A) 3.75 minutes
- B) 5 minutes**
- C) 7 minutes
- D) 10 minutes
- E) 15 minutes

2. ^{210}Bi (an isotope of bismuth) has a half-life of 5.0 days. The time for three-quarters of a sample of ^{210}Bi to decay is:

- A) 2.5 days
- B) 3.75 days
- C) 10 days**
- D) 15 days
- E) 20 days

3. Radioactive ^{90}Sr has a half-life of 30 years. What percent of a sample of ^{90}Sr will remain after 60 years?

- A) 0%
- B) 14%
- C) 25%**
- D) 50%
- E) 75%

4. The half-life of a radioactive isotope is 6.5 h. If there are initially 48×10^{32} atoms of this isotope, the number of atoms of this isotope remaining after 26 h is:

- A) 12×10^{32}
- B) 6×10^{32}
- C) 3×10^{32}**
- D) 6×10^4
- E) 3×10^2

5. At the end of 14 min, $1/16$ of a sample of radioactive polonium remains. The corresponding half-life is:

- A) $(7/8)$ min
- B) $(8/7)$ min
- C) $(7/4)$ min
- D) $(7/2)$ min**
- E) $(14/3)$ min

6. The half-life of a radioactive isotope is 140 days. In how many days does the decay rate of a sample of this isotope decrease to one fourth its initial decay rate?

- A) 35 days
- B) 70 days
- C) 105 days
- D) 210 days
- E) 280 days

7. ^{40}K decays to ^{40}Ar with a half-life of 1.25×10^9 yr. Assume that rocks contain no ^{40}Ar when they form, and that the only way ^{40}Ar can be present is through the decay of ^{40}K . If the ratio of ^{40}K to ^{40}Ar in a particular rock is found to be 1:3, what is the age of the rock?

- A) 1.25×10^9 yr
- B) 2.50×10^9 yr
- C) 3.75×10^9 yr
- D) 5.00×10^9 yr
- E) cannot be determined without knowing how much ^{40}K was in the rock to begin with

8. An isotope of Tc having a half-life of 6.0 h is used in bone scans. If a certain amount of this Tc is injected into the body, how long does it take for its initial decay rate to decrease BY 99%?

- A) 0.060 h
- B) 3.3 h
- C) 33 h
- D) 40 h
- E) slightly more than a month

9. The ratio of the radius of a classical electron (2.8×10^{-15} m) to the radius of a ^4He nucleus is

- A) 2.0
- B) 0.68
- C) 1.47
- D) 0.92
- E) 2.4

10. A certain isotope has a half-life of 32.4 hr and a relative biological effectiveness of 3.50. A sample of this isotope initially delivers an absorbed dose of 0.240 Gy to 250 g of tissue.

- (a) What was the initial equivalent dose to the tissue in rem and in Sv? **84 rem, 0.84 Sv**
- (b) What energy (in J) did the 250-g sample initially receive from the isotope? **0.06 J**

11. The maximum permissible workday dose for occupational exposure to radiation is 26 mrem. A 55-kg laboratory technician absorbs 3.3 mJ of 0.40-MeV gamma rays in a workday. The relative biological effectiveness (RBE) for gamma rays is 1.00. What is the ratio of the equivalent dosage received by the technician to the maximum permissible equivalent dosage?

- A) 0.23
- B) 0.25
- C) 0.28
- D) 0.30
- E) 0.32

12. A 70-kg laboratory technician absorbs 2.9 mJ of 0.50-MeV gamma rays in a workday. How many gamma-ray photons does the technician absorb in a workday?

- A) 3.6×10^{10}
- B) 3.6×10^9
- C) 3.6×10^8
- D) 1.0×10^9
- E) 1.0×10^8

13. A 57-kg researcher absorbs 6.3×10^8 neutrons in a workday. The energy of the neutrons is 2.6 MeV. The RBE for fast neutrons is 10. What is the equivalent dosage of the radiation exposure, in mrem, of this worker?

- A) 4.6 mrem
- B) 1.4 mrem
- C) 2.9 mrem
- D) 14 mrem
- E) 46 mrem

14. The radioactive nuclei ^{60}Co is widely used in medical applications. It undergoes beta decay, and the total energy of the decay process is 2.82 MeV per decay event. The half-life of this nucleus is 272 days. Suppose that a patient is given a dose of 6.9 μCi of ^{60}Co . If all of this material decayed while in the patient's body, what would be the total energy deposited there? (1 Ci = 3.70×10^{10} decays/s)

- A) 11 J
- B) 8.6 GJ
- C) 3.9 J
- D) 24 J
- E) 4.15 MJ

15. A laboratory experiment uses a $10\ \mu\text{Ci}\ ^{137}\text{Cs}$ source. Each decay emits a 0.66 MeV gamma ray.

- A 60 kg person standing nearby absorbs 10 % of the gamma rays. What is his absorbed dose in rads in 1 hour? $2.34 \times 10^{-5}\ \text{rad}$
- Find his effective dose in rems (take RBE = 0.8). $0.0187\ \text{mrem}$

16. Suppose your last **physical exam** included a chest X-ray, during which you received a dose of $60\ \mu\text{Sv}$.

- What was your dose in mrem? $6\ \text{mrem}$
- What was the absorbed dose in μGy and mrad? $60\ \mu\text{Gy}$, $6\ \text{mrad}$.
- How much energy did you absorb, assuming that the X-rays illuminated 15 kg of your body? $9 \times 10^{-4}\ \text{J}$

Hmm... I bet that you prefer to go through 10 of such diagnostic X-ray **physical exams** than being asked to sit for your **"105" physics exam!** Don't you?