

# Introductory to Microbiology

---

Anas Abu-  
Humaidan M.D.  
Ph.D.



## ENTEROBACTERIACEAE / Classification

G

- VE

لقد *E. coli* 11 فصلا

- More than 50 genera and hundreds of species and subspecies, Enterobacteriaceae are **ubiquitous** (سلي) organisms found worldwide in soil, water, and vegetation and are part of the normal intestinal flora of most <sup>Normal</sup> animals, including humans ).  
*Flora* *Parve* *laga*
- In humans it can be part of the **normal intestinal flora**, or <sup>دائم</sup> **always associated with human disease, or opportunistic infections**, or normally commensal organisms that <sup>virulence</sup> become pathogenic when they acquire virulence genes.

- Enterobacteriaceae are moderate-sized (0.3 to 1.0 × 1.0 to 6.0 μm), **non-spore-forming, gram-negative rods**, **facultative anaerobes** that share a common antigen **enterobacterial common antigen**

5, w

### Scientific classification

Domain:	Bacteria
Phylum:	Proteobacteria
Class:	Gammaproteobacteria
Order:	Enterobacterales
Family:	<b>Enterobacteriaceae</b>
	Rahn, 1937

Several pathogens  
بعض الممرضات



# ENTEROBACTERIACEAE / Pathogenesis and Immunity

- **Endotoxin** toxin activity depends on the **lipid A** component of LPS, released at cell lysis.
- **Capsule**, hydrophilic capsular antigens repel the hydrophobic phagocytic cell surface, but anticapsular antibodies diminish the capsule role.
- **Antigenic Phase Variation**, somatic O antigens, capsular K antigens, and flagellar H antigens alternately expressed or not expressed (phase variation).
- **Type III Secretion Systems**.  
*Secrete exotoxins*  
*Fe ١١ سرقو*  
*145 virulence expression changes in every phase*
- The bacteria counteract **iron sequestration** by producing their own competitive **siderophores** or iron-chelating compounds (e.g., enterobactin, aerobactin). Or from iron released from lysed cells.  
*145*  
*ليس نيشر بطور جوار ايكلي*
- Resistance to Serum Killing and Antimicrobial Resistance.



# ENTEROBACTERIACEAE/MacConkey's agar

Identification + purification

- It contains **bile salts** (to inhibit most Gram-positive bacteria), **crystal violet dye** (which also inhibits certain Gram-positive bacteria), **neutral red dye** (which turns pink if

selecting -ve

the microbes are fermenting

(lactose).

و في طريقه تبيّن  
للإعلاميّة

Differentiation

Lactose

Fermenters

- Enterobacteriaceae are classified based on biochemical properties, antigenic structure, and molecular analysis of their genomes, protein composition by mass spectrometry.

Lactic acid

تكون

و ينزل ال pH

و ينظر لون الطول بادي كاشف



MacConkey's agar showing

both lactose and non-lactose fermenting colonies. Lactose fermenting colonies are pink whereas non-lactose fermenting ones are colourless or appear same as the medium.

# ENTEROBACTERIACEAE/ Antibiotic resistance

لا بكتريا - لاكتام

Membrane 1  
AB 11  
2

- Resistance of the Enterobacteriaceae to antibiotics, especially of the  $\beta$  lactam type, is increasingly dominated by the mobilization of continuously expressed single genes that encode efficient drug modifying enzymes.


لا بكتريا - لاكتام

- Multi drug resistant (MDR)** Enterobacteriaceae has been frequently reported from different parts of the world as an emergence of treatment problem. Antibiotics given empirically without proper antibiotic susceptibility testing are one of the major causes for the development of MDR.

- There is a shift of the "natural" resistance, such as **membrane impermeability** and **drug efflux**, to the modern paradigm of **mobile gene pools** that transmit resistance between bacteria.

Horizontal gene transfer  
live

- <https://www.bmj.com/content/352/bmj.h6420>



## ENTEROBACTERIACEAE / overview

The following pathogens are discussed

- آ كبر مسبب لآ حرافا UT و UT بانياف
- *Escherichia coli* { كلها لعلوا
  - *Salmonella* ✓
  - *Shigella* ✓ diarrhea
  - *Yersinia* → plague { بار
  - *Klebsiella* G1
  - *Proteus* → UTI
- Neisseria meningitidis, G1, UTI
- Staphylococcus aureus
- Staph + strep



## -


- *E. coli* is the most common and important member of the genus *Escherichia*.

- It is both a common **commensal** inhabitant of the

# pathogens in humans.

- It is a frequent cause of diarrheal disease.

- 1) 1/2  
2) 1/2



## *Escherichia coli* strains <sup>3) B stream</sup>

لدينا كميات كبيرة لخمير

- **Commensal** strains innocuously colonize the colon of healthy hosts , causing extraintestinal disease only in the presence of a large inoculum (e.g., with penetrating abdominal trauma) and/or significant host compromise.
- **Diarrhoeagenic strains cause diarrhoea syndromes** that vary in clinical presentation and pathogenesis according to the strain's distinctive virulence traits
- **Extraintestinal pathogenic *E. coli*** (ExPEC) often innocuously colonize the human gut. However, they have a unique ability to enter and survive within normally sterile extraintestinal body sites, and to **cause disease** when they do so.

يمكن تلخيصها بـ 3 أنواع على الأقل هي: *infection* و *كائنات* و *تسبب المرض*

Organism	Site of Action	Disease	Pathogenesis
Enterotoxigenic <i>E. coli</i> (ETEC)	Small intestine	Traveler's diarrhea; infant diarrhea in developing countries; watery diarrhea, vomiting, cramps, nausea, low-grade fever	Plasmid-mediated, heat-stable (ST) and heat-labile (LT) enterotoxins that stimulate hypersecretion of fluids and electrolytes
Enteropathogenic <i>E. coli</i> (EPEC)	Small intestine	Infant diarrhea in developing countries; watery diarrhea and vomiting, nonbloody stools; believed to be rare in United States	Plasmid-mediated A/E histopathology, with disruption of normal microvillus structure resulting in malabsorption and diarrhea
Enteraggregative <i>E. coli</i> (EAEC)	Small intestine	Infant diarrhea in developing and probably developed countries; traveler's diarrhea; persistent watery diarrhea with vomiting, dehydration, and low-grade fever	Plasmid-mediated aggregative adherence of rods ("stacked bricks") with shortening of microvilli, mononuclear infiltration, and hemorrhage; decreased fluid absorption
Shiga toxin-producing <i>E. coli</i> (STEC)	Large intestine	Initial watery diarrhea followed by grossly bloody diarrhea (hemorrhagic colitis) with abdominal cramps; little or no fever; may progress to hemolytic uremic syndrome	STEC evolved from EPEC; A/E lesions with destruction of intestinal microvilli, resulting in decreased absorption; pathology mediated by cytotoxic Shiga toxins (Stx1, Stx2), which disrupt protein synthesis
Enteroinvasive <i>E. coli</i> (EIEC)	Large intestine	Rare in developing and developed countries; fever, cramping, watery diarrhea; may progress to dysentery with scant bloody stools	Plasmid-mediated invasion and destruction of epithelial cells lining colon

Stool  
Watery  
Fecal contaminated

# Enterotoxigenic E. coli (ETEC)

- one of the most common causes of bacterial diarrheal disease in developing countries, and 30% of traveler's diarrhea\*. Acquired through consumption of fecally contaminated food or water. Person-to-person spread does not occur.
- 1- to 2-day incubation period and persists for an average of 3 to 5 days.

- The symptoms: **Secretory diarrhea** (watery, non-bloody diarrhea) and **abdominal cramps**; less commonly nausea and vomiting. Can be fatal in undernourished individuals.
  - Produces 2 classes of toxins:
- Feco - GABA route

Heat stable toxin leads to **increase in cyclic guanosine monophosphate (cGMP)** and subsequent hypersecretion of fluids well as inhibition of fluid absorption

heat labile toxins leads to **increase in cyclic adenosine monophosphate (cAMP)** levels, resulting in enhanced secretion of chloride and decreased absorption of sodium and chloride

**\*traveler's diarrhea :** When you visit a place where the climate or sanitary practices are different from yours at home, you have an increased risk of developing traveler's diarrhea. Usually recovery happens within days with no need for treatment.



## Enterotoxigenic *E. coli* (ETEC)

A second-year medical student experiences **watery diarrhea and mild abdominal cramps** during his 2-week travel to Egypt. With his little medical knowledge, he makes several assumptions, which of those assumption is **false**?

- a) This is probably a case of traveler's diarrhea that should resolve within a few days. ✓
- b) Enterotoxigenic *E. coli* (ETEC) is a probable causative agent. ✓
- c) He would not have become sick if he washed his hands properly.
- d) Liquids are important to prevent dehydration and loss of electrolytes.
- e) If it is traveler's diarrhea, he probably contracted the pathogen in a meal he ate 2 days ago.

(Bloody) Mostly watery  
أغلبها سائل

watery diarrhea  
سائل  
أغلبه سائل  
في يومين أو ثلاثة  
في يومين أو ثلاثة

## Shiga toxin-producing E. coli (STEC)

ع. coli  
بكتريا شيجا  
1 خلية (تؤكل)

فوكي  
Virulence  
Factor  
بهره من انا

- Most infections are attributed to the consumption of undercooked meat products, water, unpasteurized milk or fruit juices, uncooked vegetables, and fruits. Ingestion of fewer than 100 bacteria can produce disease, and person-to-person spread occurs.

low inoculum

colony  
عده  
الواحد في انا

- Disease caused by STEC ranges from mild uncomplicated diarrhea to hemorrhagic colitis with severe abdominal pain and bloody diarrhea. Severe disease is more commonly associated with STEC O157:H7.

- 3 to 4 days of incubation, Within 2 days of onset, disease in 30% to 65% of patients progresses to a bloody diarrhea with severe abdominal pain, Complete resolution of symptoms typically occurs after 4 to 10 days in most untreated patients.

HUS

Endothelial  
بكتريا انا  
ABc و انا

- Hemolytic uremic syndrome (HUS)**, a disorder characterized by acute renal failure, thrombocytopenia, and microangiopathic hemolytic anemia, is a complication in 5% to 10% of infected children younger than 10 years.

نفسه  
نفسه  
نفسه

Shiga toxin

Cell wall  
Antigen  
Hsp  
Secretory



## Clinical Case 25-1 Multistate Outbreak of Shiga Toxin-Producing *Escherichia coli* (STEC) Infections

In 2006, *E. coli* O157 was responsible for a large multistate outbreak of gastroenteritis. The outbreak was linked to contamination of spinach, with a total of 173 cases reported in 25 states, primarily over an 18-day period. The outbreak resulted in hospitalization of more than 50% of the patients with documented disease, a 16% rate of hemolytic uremic syndrome, and one death. Despite the wide distribution of the contaminated spinach, publication of the outbreak and the rapid determination that spinach was responsible resulted in prompt removal of spinach from grocery stores and termination of the outbreak. This outbreak illustrates how contamination of a food product, even with small numbers of organisms, can lead to a widespread outbreak with a particularly virulent organism, such as strains of STEC.

Shiga-Toxin *E. coli* Hemolytic Uremic Syndrome: Review of Management and Long-term Outcome

<https://link.springer.com/article/10.1007/s40124-020-00208-7>

1) UTI  
3) septicemia

# Escherichia coli/ Extraintestinal Infections

Meningitis

بكتيريا سالبة الجرام

80%

- **Urinary Tract Infection:** Most gram-negative rods that produce UTIs originate in the colon, contaminate the urethra, ascend into the bladder, and may migrate to the kidney, prostate,   
 catheter

- Almost every second woman suffers from a bladder infection at some point in her life ( **E. coli in 80% of UTI cases** ). Also men are affected by cystitis, though less frequently, probably due to anatomical differences (e.g. shorter urethra in women makes it easier for bacteria to reach the bladder).
- **Neonatal Meningitis** : *E. coli* and group B streptococci cause the majority of CNS infections in infants younger than 1 month.

Birth Canal Baby

1) *Colitis*

2

3) *Monocytogens*

True

-ve → *Septicemia*


sepsis

- **Septicemia** : Typically, septicemia caused by gram-negative rods, such as *E. coli*, most commonly originates from infections in the urinary or GI tract, with high mortality in immunocompromised patients.



*Escherichia coli*

## Extraintestinal Infections



# Salmonella

Normal  
Alivra  
ميكروبيوتا

سلالة بكتيريا خارج الخلية  
دوائية

- Salmonella can colonize virtually all animals (especially poultry). Serotypes such as Salmonella Typhi and Salmonella Paratyphi are highly adapted to humans and do not cause disease in nonhuman hosts.

Human  
ميكروبيوتا  
Mucosal

- After ingestion and passage through the stomach, salmonellae attach to the mucosa of the small intestine and invade into the M (microfold) cells located in Peyer patches, as well as into enterocytes. The bacteria remain in endocytic vacuoles, where they replicate. The bacteria can also be transported across the cytoplasm and released into the blood or lymphatic circulation. The inflammatory response confines the infection to the GI tract, mediates the release of prostaglandins, and stimulates cAMP and active fluid secretion.  
Virulence  
آلية في الخلية التي تسمى

- Virulence dependent on pathogenicity island on the bacterial chromosome. Encoding for toxins, attachment proteins and immune evasion mechanisms.

Salmonella  
E.C  
Intra cellularly  
Salmonella  
البكتيريا  
+  
Salmonella  
سلالة بكتيريا

به دوتا آكله

جوانا اسامه  
Salmonella

وهيك بروجع على ال BV وينقل

به دوا آكله

جوانا الجسم

### • Asymptomatic Colonization :

The strains of *Salmonella* responsible for causing typhoid and paratyphoid fevers are maintained by human colonization.



## Mary Mallon

Cook

Mary Mallon, also known as **Typhoid Mary**, was an Irish cook. She was the first person in the United States identified as an asymptomatic carrier of the pathogen associated with typhoid fever. She was presumed to have infected 51 people, three of whom died, over the course of her career as a cook. [Wikipedia](#)

**Born:** September 23, 1869, Cookstown, United Kingdom

**Died:** November 11, 1938, Riverside Hospital

# Salmonella / Epidemiology and diseases

- The most common sources of human infections are **poultry, eggs, dairy products**, and foods prepared on contaminated work surfaces, **large inoculum** (e.g.,  $10^6$  to  $10^8$  bacteria) is required for symptomatic disease.  
(في الـ ع.ع. العدائية) (أما الـ Shiga و E. coli السريع)
- The infectious dose for *Salmonella* Typhi infections is **low**, so **person-to-person spread is common**, occur when food or water contaminated by infected food handlers is ingested.  
من الـ جعة و Next bloody
- **Gastroenteritis** is a **common form of salmonellosis**, nausea, vomiting, and nonbloody diarrhea. can persist for 2 to 7 days before spontaneous resolution.  
في الـ Traveler's E. coli و ذلك الـ شللي
- **Septicemia** All *Salmonella* species can cause bacteremia, although infections with *Salmonella* Typhi, *Salmonella* Paratyphi more commonly lead to a bacteremic phase.

في الـ الـ خطر

1) G-Iitis  
2) septicemia




# Salmonella / diseases

التي ينتجها M (مادة) التي تفرغها

- *Salmonella* Typhi produces a febrile illness called **typhoid fever**. A milder form of this disease, referred to as **paratyphoid fever**, is produced by other *Salmonella* (e.g. *paratyphi*).

- The bacteria responsible for enteric fever pass through the cells lining the intestines and are engulfed by macrophages. They replicate after being transported to the liver, spleen, and bone marrow. Ten to 14 days after ingestion of the bacteria, patients experience gradually increasing fever, with nonspecific complaints of headache, myalgias, malaise, and anorexia).

في اللمفاوية



# Salmonella / diseases




## Clinical Case 25-2 *Salmonella* Typhi Infection

Scully and associates (*N Engl J Med* 345:201–205, 2007) described a 25-year-old woman who was admitted to a Boston hospital with a history of persistent fever that did not respond to amoxicillin or acetaminophen or ibuprofen. She was a resident of the Philippines who had been traveling in the United States for the previous 11 days. On physical examination, she was febrile and had an enlarged liver, abdominal pain, and an abnormal urinalysis. Blood cultures were collected upon admission to the hospital and were positive the next day with *Salmonella* Typhi. Because the organism was susceptible to fluoroquinolones, this therapy was selected. Within 4 days, she had defervesced and was discharged to return home to the Philippines. Although typhoid fever can be a very serious life-threatening illness, it can initially present with nonspecific symptoms, as was seen in this woman.

### Typhoid Fever

<https://www.nejm.org/doi/full/10.1056/nejmra020201>



# Shigella


شيجلا  
التي تسبب التهاب القولون البكتيري

- *S. dysenteriae*, *Shigella flexneri*, *Shigella boydii*, and *Shigella sonnei*. However, analysis of DNA has determined that these four species are actually biogroups within the species *E. coli*.
- *Shigellae* cause disease by invading and replicating in cells lining the **colon**. Structural gene proteins mediate the adherence of the organisms to the cells, as well as their invasion, intracellular replication, and cell-to-cell spread.
- *S. dysenteriae* strains produce an exotoxin, **Shiga toxin**. Similar to Shiga toxin produced by **STEC**
- The A subunit in the toxin cleaves the **28S rRNA** in the 60S ribosomal subunit, thereby preventing the binding of aminoacyl-transfer RNA and **disrupting protein synthesis**.
- The primary manifestation of toxin activity is **damage to the intestinal epithelium**; however, in a small subset of patients, the Shiga toxin can mediate **damage to the glomerular endothelial cells, resulting in renal failure (HUS)**.

severe type of E.C

## Shigella - الـ شيجيلا

Shigella passes the epithelial cell (EC) barrier by transcytosis through M cells and encounters resident macrophages. The bacteria evade degradation in macrophages by inducing an apoptosis-like cell death, which is accompanied by proinflammatory signaling. Free bacteria invade the EC from the basolateral side, move into the cytoplasm by actin polymerization, and spread to adjacent cells. Proinflammatory signaling by macrophages and EC further activates the innate immune response and attracts PMN. The influx of PMN disintegrates the EC lining, which initially exacerbates the infection and tissue destruction by facilitating the invasion of more bacteria. Ultimately, PMN phagocytose and kill Shigella, thus contributing to the resolution of the infection.



# Shigella / Epidemiology and diseases

*Sonnei* → U.S

*flexneri*

*dysenteriae* → West Afr. + Ame.

- Humans are the only reservoir for *Shigella*.
- *S. sonnei* is responsible for almost 85% of U.S. infections, whereas *S. flexneri* predominates in developing countries. Epidemics of *S. dysenteriae* infections occur periodically, most recently in West Africa and Central America.

- *Shigellosis* (*Shigella* infection) ( is primarily a pediatric disease, with 60% of all infections in children younger than 10 years.

Person to person low incubation

- *Shigellosis* is transmitted person to person by the fecal-oral route. Because as few as 100 to 200 bacteria can establish disease, shigellosis spreads rapidly in communities where sanitary standards and the level of personal hygiene are low.

STEC 57

- *Shigellosis* is characterized by abdominal cramps, diarrhea, fever, and bloody stools. The clinical signs and symptoms of the disease appear 1 to 3 days after the bacteria are ingested.

- Infection is generally self-limited, although antibiotic treatment is recommended to reduce the risk of secondary spread to family members and other contacts.

Salmonella



Shigella



## Clinical Case 25-3 *Shigella* Infections in Day-Care Centers

In 2005, three states reported outbreaks of multidrug-resistant *Shigella* infections in day-care centers. A total of 532 infections were reported in the Kansas City area, with the median age of patients 6 years old (Centers for Disease Control and Prevention: *MMWR Morb Mortal Wkly Rep* 55:1068–1071, 2006). The predominant pathogen was a multidrug-resistant strain of *Shigella sonnei*, with 89% of the isolates resistant to ampicillin and trimethoprim-sulfamethoxazole. Shigellosis spreads easily in day-care centers because of the increased risk of fecal contamination and the low infectious dose responsible for disease. Parents and teachers, as well as classmates, are at significant risk for disease.

(\*)  
ref

E.C. 11a Quesi nA 11

# Klebsiella

Nose,  
Mouth,  
الوجنة

- Klebsiella species are routinely found in the human nose, mouth, and gastrointestinal tract as normal flora. *فلاورا* *فوزية* *نورس* *نورس* *Nosocomial*
- The most commonly isolated members of this genus are *K. pneumoniae*, which can cause community- or hospital-acquired primary lobar pneumonia. These bacteria also cause wound and soft-tissue infections and UTIs.
- The ability of *K. pneumoniae* to colonize the hospital environment, including carpeting, sinks, flowers, and various surfaces, as well as the skin of patients and hospital staff, has been identified as a major factor in the spread of hospital-acquired infections.

\*



## Proteus


UTI

urease

PH ٨  
بشكل جيد  
احضار بنى ربي  
بشكل طبيعي

- *P. mirabilis*, the most common member of this genus, primarily produces **infections of the urinary tract**.

- *P. mirabilis* produces large quantities of **urease**, which splits **urea into carbon dioxide and ammonia**. This process raises the urine pH, precipitating magnesium and calcium in the form of **struvite and apatite crystals**, respectively, and results in the formation of **renal (kidney) stones**. The increased alkalinity of the urine is also toxic to the uroepithelium



لحمية  
بشكل جيد  
Stones

Pestis

بشكل طبيعي

***Yersinia***

Hispanic / path  
الطائف

Mortality Rate of  
% 57

- The best-known human pathogen within the genus *Yersinia* is *Y. pestis* جاتي من الصوفان
- All *Yersinia* infections are **zoonotic**, with humans the accidental hosts. There are two forms of *Y. pestis* infection: **urban plague**, for which rats are the natural reservoirs, and **sylvatic plague**, which causes infections in squirrels, rabbits, field rats, and domestic cats.
- **3 major pandemics that shaped history.**

Urban

Sylvatic

Bubonic


Rats reservoir

# Yersinia <sup>تسمى</sup> الطاعون <sup>أو</sup> كس الحرق <sup>أو</sup> كس الحرق <sup>أو</sup> كس الحرق

- Bubonic plague caused by *Y. pestis* is characterized by an incubation period of no more than 7 days after a person has been bitten by an infected flea. Patients have a high fever and a painful **bubo** (inflammatory swelling of the lymph nodes) in the groin or axilla. Bacteremia develops rapidly if patients are not treated, and as many as 75% die. <sup>أول على الفئران يهون</sup>

- The patients are highly infectious; person-to-person spread occurs by aerosols in case of **pneumonic plague**. <sup>و يمكن به ك</sup>

أول كس الحرق





الناس



ما تضرها ٥٠ ألفا


The Justinian Plague began in 541 AD and was followed by frequent outbreaks over the next two hundred years that

eventually killed over 25 million people and affected much of the Mediterranean basin—virtually all of the known world at that time.

الرومان

عالمية ربه ان الله عليه اجعت

الدكتورة



Mask

كانوا بذكروا

الاهل

بنتقل من

الاهل



حيروا انه ينتقل بالقران


The third pandemic, the Modern Plague, began in China in the 1860s and appeared in Hong Kong by 1894. Over the next 20 years, it spread to port cities around the world by rats on steamships. The pandemic caused approximately 10 million deaths

انتشر

The second pandemic, widely known as the "Black Death" or the Great Plague, originated in China in 1334 and spread along the great trade routes to Constantinople and then to Europe, where it claimed an estimated 60% of the European population, around 50-200 million lives.

Europe ۛ

## The evolution of bacteria on a "mega-plate" Petri dish



## **Further reading:**

- Murray - Medical Microbiology 8th Edition  
Section 4: Bacteriology  
Chapter 25: ENTEROBACTERIACEAE