


# Reviewing the Literature

## Chapter 3

تحت اهيها في تقرير ادوية  
التي اهيها في تقرير ادوية  
التي اهيها في تقرير ادوية  
التي اهيها في تقرير ادوية  
more  
refinement  
للدراسة



teams.microsoft.com is sharing your screen.

Stop sharing

Hide

© 2014 by Jones & Bartlett Learning, LLC an Ascend Learning Company  
www.jblearning.com

معلومات  
غير رسمية

3

### 3.1 Informal Sources

- Nontechnical information from trusted sources (e.g., CDC, WHO) can provide helpful background on a topic.
- Factsheets and other informal information are not part of the formal peer-reviewed scientific literature (Do not cite them in formal reports).

معلومات غير رسمية  
الأمم المتحدة  
(Declaration)  
التي هو صواب  
التي دلالة  
بمعرفة

## 3.2 Statistical Reports

formel  
an die  
UN, in  
Referenz

- World Bank world development indicators
- UN agency reports (*World Health Statistics, Human Development Report, State of the World's Children*)
- Annual reports from groups like the American Cancer Society and Population Reference Bureau
- Information from state and local health departments

المقالة  
البيانية

# 3.3 Abstract Databases (1 of 4)

- **Abstract:** a paragraph-length summary of an article, chapter, or book.
- **Structured Vs. Unstructured.**
- Use **keywords** to search multiple abstract databases.

Brief Description of study Designs, Findings, Results.

SUBSTANCE USE & MISUSE  
https://doi.org/10.1080/10826084.2019.1645177

ORIGINAL ARTICLE

**Waterpipe Tobacco Smoking among University Students in Three Eastern Mediterranean Countries: Patterns, Place, and Price**

Ramzi G. Salloum<sup>a,b</sup>, JuHan Lee<sup>a</sup>, Aya Mostafa<sup>c</sup>, Niveen M.E. Abu-Rmeileh<sup>d</sup>, Randah R. Hamadeh<sup>e</sup>, Muhammad W. Darawad<sup>f</sup>, Khalid A. Kheirallah<sup>g</sup>, Mohamed Salama<sup>h</sup>, Wasim Maziak<sup>i</sup>, and Rima Nakkash<sup>b</sup>

<sup>a</sup>Department of Health Outcomes and Biomedical Informatics, College of Medicine, University of Florida, Gainesville, Florida, USA; <sup>b</sup>Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon; <sup>c</sup>Faculty of Medicine, Ain Shams University, Cairo, Egypt; <sup>d</sup>Institute of Community and Public Health, Birzeit University, Birzeit Palestine; <sup>e</sup>College of Medicine and Medical Sciences, Arabian Gulf University, Manama, Bahrain; <sup>f</sup>School of Nursing, The University of Jordan, Amman, Jordan; <sup>g</sup>Faculty of Medicine, Jordan University of Science and Technology, Irbid, Jordan; <sup>h</sup>Medical Experimental Research Center, Mansoura University, Mansoura, Egypt; <sup>i</sup>Department of Epidemiology, Florida International University, Miami, Florida, USA

**ABSTRACT**  
**Background:** The objective of this study was to examine waterpipe tobacco smoking patterns, places of smoking, and prices paid among university students in the Eastern Mediterranean Region. **Methods:** A cross-sectional online survey was administered to a convenience sample of university students in three countries. Participants were young adults (18–29 years) who were ever waterpipe smokers, from Egypt (n = 728), Jordan (n = 790), and Palestine (n = 722). Measures included past-30-day waterpipe smoking, frequency, intensity, place of smoking, and prices paid per waterpipe smoking session and for packaged waterpipe tobacco. Logistic regression models evaluated the factors associated with past-30-day waterpipe smoking. **Results:** Past-30-day waterpipe smoking (prevalence) was observed among 60.7%, 67.7% and 63.1% of students from Egypt, Jordan, and Palestine, respectively. Among past-30-day smokers, past-5-day waterpipe smoking (frequency) was observed among 28.9%, 51.5%, and 48.6% of participants, respectively. Smoking in a café was highest among participants from Egypt (74.0%), followed by those from Palestine (44.8%), and Jordan (43.0%). Mean price paid per session was USD 0.99 (Egypt), USD 8.07 (Jordan), USD 6.05 (Palestine). The corresponding mean prices per packet were USD 0.86, USD 4.96, and USD 5.55, respectively. Predictors of past-30-day waterpipe smoking included younger age of initiation, male gender, employment, and smoking waterpipe alone. **Conclusions:** This study contributes to our understanding of waterpipe smoking patterns, places of smoking, and prices paid among young adults in a region with alarmingly high smoking rates. Understanding waterpipe smoking behaviors can inform the design of policy and educational interventions to curb its rising threat.

**KEYWORDS**  
 Hookah, waterpipe; shisha; smoking; Eastern Mediterranean

المقالة  
البيانية

# Unstructured Abstract

عنوان  
مستخلص  
النتائج

## Evaluating the Psychometric Properties of the Arabic Version of the Diabetes Distress Scale

Muhammad W. Darawad, PhD, RN; Sawsan Hammad, RN, MSN; Osama A. Samarkandi, PhD, RN; Ayman M. Hamdan-Mansour, PhD, RN; and Amani A. Khalil, PhD, RN

### ABSTRACT

The current study evaluated the psychometric properties of the Arabic version of the Diabetes Distress Scale (DDS-A) among Arab patients with diabetes mellitus (DM) using a descriptive cross-sectional design. Participants' DDS-A total scores significantly correlated with depressive symptoms ( $r = 0.288, p = 0.00$  [Query #1:  $p$  value cannot equal 0.00. Please change to  $p < 0.05, p < 0.01, etc.,$  throughout the abstract and article text.]) as well as two subscales, emotional

burden ( $r = 0.276, p = 0.00$ ) and regimen distress ( $r = 0.265, p = 0.00$ ). Participants' DDS-A scores had significant negative correlations with income, DM self-management, and knowledge ( $r = -0.184, p = 0.008; r = -0.310, p = 0.00; r = -0.174, p = 0.003$ , respectively) and a positive correlation with HbA1C level ( $r = 0.153, p = 0.018$ ). Factor analysis revealed a four-factor solution that retained all items and explained a variance of 65.59%. Cronbach's alpha was 0.822 for the total scale and 0.778 to 0.881 for the sub-

scales, indicating a high internal consistency. The DDS-A was found to be a psychometrically sound measure to evaluate DM-related distress among Arab patients. [*Journal of Psychosocial Nursing and Mental Health Services*, xx(x), xx-xx.]

Diabetes mellitus (DM) is a chronic disease that affects 366 million individuals worldwide, a number that is expected to rise by 2030 to 552 million (Whiting, Guariguata, Weil, & Shaw, 2011). Developing countries are at a higher risk for more DM cases, with a 69% increase in the number of adults with

## 3.3 Abstract Databases (2 of 4)

Examples of databases that are free to the public:

- PubMed
- European PubMed Central (PMC)
- SciELO & LILACS (Central and South America)
- AJOL (Africa)

هذه مواقع تقيض  
Abstract  
تقيض (أي) ملخص  
بحيث وقت ينور على  
الموضوع ليس كذا  
→ Abstract

## 3.3 Abstract Databases (3 of 4)

Examples of other (usually with subscription) databases:

- MEDLINE
- CINAHL
- Embase
- PsycINFO
- Web of Science
- EBSCO, JSTOR, Ovid, and ProQuest
- Company-specific databases (LWW, SAGE, T&F, Wiley, others)

نیشنل ایس ایم  
Abstract کی  
مطابقت  
اعتماد  
کا عملہ





## 3.4 Full-Text Articles

### Where to find free full-text PDFs:

- Google Scholar and other search engines
- PubMed Central and other open access repositories
- Journal websites (if the article is open access)
- Library subscriptions (e-journals) or interlibrary loans when a journal is not in a library's collection
- E-mail the author to politely request a copy
- <http://e-library> (inside JU)

لکھو  
لکھو  
)

teams.microsoft.com is sharing your screen. Stop sharing Hide

Jones & Bartlett Learning, LLC an Ascend Learning Company  
www.jblearning.com

# 3.5 Critical Reading

فصل ٣  
الدراسة

- هل تم إجراء الدراسة بشكل صحيح ؟  
- هل أداة القياس مناسبة ؟  
- هل النتائج صحيحة ؟  
- هل الاستنتاج صحيح ؟

- **Internal validity:** How well was the study designed, conducted, interpreted, and reported?
- **External validity (generalizability):** How likely is it that the results of this study apply to other populations?

Internal validity  
الصدق الداخلي  
: هو موثوق نسبي  
أي مدى  
معيّن وليس  
أبديتها وأسود

نتائج الدراسة هل يمكن تعميمها على  
آخرين