## Cardiac output and Venous Return


Faisal I. Mohammed, MD, PhD

## Objectives

- Define cardiac output and venous return
- Describe the methods of measurement of CO
- Outline the factors that regulate cardiac output
- Follow up the cardiac output curves at different physiological states
- Define venous return and describe venous return curve
- Outline the factors that regulate venous return curve at different physiological states
- Inter-relate Cardiac output and venous return curves


# **Important Concepts About Cardiac Output (CO) Control**

- Cardiac Output is the sum of all tissue flows and is affected by their regulation (CO = 5L/min, cardiac index = 3L/min/m<sup>2</sup>).
- CO is proportional to tissue  $O_2$  use.
- CO is proportional to 1/TPR when AP is constant.
- $F = \Delta P/R$  (Ohm's law)
- CO = (MAP RAP) / TPR, (RAP=0) then
- CO=MAP/TPR; MAP=CO\*TPR



WORK OUTPUT DURING EXERCISE (kg\*m/min)


# Magnitude & Distribution of CO at Rest & During Moderate Exercise




## Variations in Tissue Blood Flow

			ml/min/
	Per cent	ml/min	100 gm
Brain	14	700	50
Heart	4	200	<b>70</b>
Bronchi	2	100	<b>25</b>
Kidneys	22	1100	360
Liver	<b>27</b>	1350	<b>95</b>
Portal	(21)	(1050)	
Arterial	(6)	(300)	
<b>Muscle (inactive state)</b>		<b>750</b>	4
Bone	5	<b>250</b>	3 3
Skin (cool weather)	6	<b>300</b>	3
Thyroid gland	1	<b>50</b>	<b>160</b>
Adrenal glands	0.5	25	<b>300</b>
Other tissues	3.5	175	1.3
Total	100.0	5000	

## Control of Cardiac Output



### Factors that affect the Cardiac Output



#### **Autonomic Effects on Heart**

- □ Sympathetic stimulation causes increased HR and increased contractility with
- HR = 180-200 and C.O. = 15-20 L/min.
- □ Parasympathetic stimulation decreases HR markedly and decreases cardiac contractility slightly. Vagal fibers go mainly to atria.
- □ Fast heart rate (tachycardia) can decrease C.O. because there is not enough time for heart to fill during diastole.


## Cardiac Contractility

- Best is to measure the C.O. curve, but this is nearly impossible in humans.
- dP/dt is not an accurate measure because this increases with increasing preload and afterload.
- (dP/dt)/P <sub>ventricle</sub> is better. P <sub>ventricle</sub> is instantaneous ventricular pressure.
- Excess K<sup>+</sup> decreases contractility.
- Excess Ca<sup>++</sup> causes spastic contraction,
   and low Ca<sup>++</sup> causes cardiac dilation.


#### Work of the ventricles

- During the latter part of the ejection phase how can blood still leave the ventricle if pressure is higher in the aorta? Momentum of blood flow
- Total energy of blood = P + mV<sup>2</sup>/2
   = pressure + kinetic energy
- Total energy of blood leaving ventricle is greater than in aorta.


#### Ventricular Stroke Work Output




Left Atrial Mean Pressure (mm Hg)




Right Atrial Mean Pressure (mm Hg)



#### **Effect of Sympathetic and Parasympathetic Stimulation on Cardiac Output**





RIGHT ATRIAL PRESSURE (mmHg)


### The Cardiac Output Curve

- Plateau of CO curve determined by heart strength (contractility + HR)
- Sympathetics  $\Rightarrow$  plateau
- $\downarrow$  Parasympathetics (HR)  $\Rightarrow$  (? plateau)
- Plateau
- Heart hypertrophy's ⇒ plateau
- Myocardial infarction  $\Rightarrow$  (? plateau)
- ↓ Plateau


#### The Cardiac Output Curve (cont'd)

- Valvular disease ⇒ ↓ plateau (stenosis or regurgitation)
- Myocarditis  $\Rightarrow \downarrow$  plateau
- Cardiac tamponade  $\Rightarrow$  (? plateau)
- ↓ Plateau
- Metabolic damage ⇒ ↓ plateau


## Factors Affecting Cardiac Output




## Factors Affecting Stroke Volume




# A Summary of the Factors Affecting Cardiac Output



#### REGULATION OF STROKE VOLUME: PRELOAD




## REGULATION OF STROKE VOLUME: CONTRACTILITY



## Cardiac Contractility

- Best is to measure the C.O. curve, but this is nearly impossible in humans.
- dP/dt is not an accurate measure because this increases with increasing preload and afterload.
- (dP/dt)/P <sub>ventricle</sub> is better. P <sub>ventricle</sub> is instantaneous ventricular pressure.
- Excess K<sup>+</sup> decreases contractility.
- Excess Ca<sup>++</sup> causes spastic contraction, and low Ca<sup>++</sup> causes cardiac dilation.

## REGULATION OF STROKE VOLUME: AFTERLOAD




## Thank You




#### Measurement of Cardiac Output

- Electromagnetic flowmeter
- Indicator dilution (dye such as cardiogreen)
- Thermal dilution
- Oxygen Fick Method
- $CO = (O_2 consumption / (A-V O_2 difference)$


## Electromagnetic flowmeter






 $q_1 = CQ^*C_{VO2}$   $q_2 = amount of Oxygen uptake by the lungs$   $q_3 = CO^* C_{AO2} \text{ and equals} = CQ^*C_{VO2} + O_2 \text{ uptake}$   $Oxygen uptake = CQ\{C_{AO2} - C_{VO2}\}$   $CO = Oxygen uptake / \{C_{AO2} - C_{VO2}\}$ 

## Spirometer




#### Swan-Ganz catheter




## O<sub>2</sub> Fick Problem

- If pulmonary vein  $O_2$  content = 200 ml  $O_2/L$  blood
- Pulmonary artery  $O_2$  content = 160 ml  $O_2$  /L blood
- Lungs add 400 ml O<sub>2</sub> /min
- What is cardiac output?
- Answer: 400/(200-160) = 10 L/min



Cardiac output = 
$$\frac{q}{C}X\frac{60}{\text{durationin seconds}}$$


#### Thermodilution Method Curve




#### VENOUS RETURN

- Definition: Volume of blood returns to either the left side or right side of the heart per minute
- $VR = CO = \Delta P/R$
- VR = (Venous pressure –Rt. Atrial pressure)/ resistance to venous return


### Effect of Venous Valves




#### Effect of Venous Valves




## Venous Valves



## Effect Of Gravity on Venous Pressure




## Vessel Structure and Function





Dilated and twisted appearance of varicose veins in the leg

## Venous Pressure in the Body




- Compressional factors tend to cause resistance to flow in large peripheral veins.
- Increases in right atrial pressure causes blood to back up into the venous system thereby increasing venous pressures.
- Abdominal pressures tend to increase venous pressures in the legs.

#### Central Venous Pressure


- \* Pressure in the right atrium is called *central venous pressure*.
- \* Right atrial pressure is determined by the balance of the heart pumping blood out of the right atrium and flow of blood from the large veins into the right atrium.
- \* Central venous pressure is normally 0 mmHg, but can be as high as 20-30 mmHg.

# Factors affecting Central Venous Pressure


- Right atrial pressure (RAP) is regulated by a balance between the ability of the heart to pump blood out of the atrium and the rate of blood flowing into the atrium from peripheral veins.
  - Factors that increase RAP:
    - -increased blood volume
    - -increased venous tone
    - dilation of arterioles
    - -decreased cardiac function




#### Factors that Facilitate Venous Return



#### The Venous Return Curve





RIGHT ATRIAL PRESSURE (mmHg)

## Venous Return (VR)

- Beriberi thiamine deficiency  $\Rightarrow$  arteriolar dilatation  $\Rightarrow \downarrow RVR$
- (RVR= resistance to venous return)
  because VR = (MSFP RAP) /RVR
  (good for positive RAP's)
- A-V fistula  $\Rightarrow$  (? RVR)
- ↓ RVR
- C. Hyperthyroidism  $\Rightarrow$  (? RVR)

## Venous Return (VR) (cont'd)

- Anemia  $\Rightarrow \downarrow RVR \text{ (why?)}$
- Sympathetics  $\Rightarrow$  MSFP
- Blood volume  $\Rightarrow$  MSFP + small
  - ↓ in RVR
- Venous compliance (muscle contraction or venous constriction)
  - $\Rightarrow$  (? MSFP)
- MSFP

## Factors Causing Venous Return

- $\downarrow$  Blood volume  $\Rightarrow \downarrow$  MSFP
- $\downarrow$  Sympathetics  $\Rightarrow$  (? v. comp. and MSFP)
- Venous compliance and ↓MSFP
- Obstruction of veins  $\Rightarrow$  (? RVR)
- RVR

RIGHT ATRIAL PRESSURE (mmHg)

## Thank You

