

Miscellaneous respiratory tract infections

By : Nader Alaridah MD,PhD

Atypical Pneumonia

- Atypical pneumonia caused by *Mycoplasma* and *Chlamydia*, *Legionella*.. These related to Gram-ve bacteria.. Attached to respiratory mucosa..Not common part of Respiratory flora..Opportunistic pathogens
- Causing mostly milder forms of pneumonia .. characterized by slow development of symptoms unlike other forms of pneumonia which can develop more quickly .. more severe early symptoms.
- *M. pneumoniae* : The smallest size Bacteria ..Lack Cell Wall.. Lipid bi-layer Membrane.. Aerobic Growth, Respiratory /Urinary Mucosa.. Various *Mycoplasma spp.* Associated with disease.. Human, Animals, Birds


Mycoplasma

- *M. pneumoniae* ..spread by droplet infection.. often develop Low fever & dry cough symptoms ..few days-weeks.. anemia, rashes, neurological syndromes..meningitis, encephalitis.
- Acute/ Subacute Pharyngitis.. Bronchitis.. Common Infection in Fall-Winter.. Mostly Old children & young Adults.
- Severe forms of M pneumonia have been described in all age groups.
- Lab Diagnosis: Special culture medium.. PCR.., Pleural fluid, Blood. Serological Cold-Agglutination Test.. Increased antibody titers.
- Treatment: levofloxacin, moxifloxacin, Macrolides/ Azithromycin.. No Vaccine

Chlamydia species

- **Chlamydia.. Attached** human mucosal membrane.. ..obligate intracellular. intracytoplasmic inclusions..Rapidly killed outside body, dryness & high temperature > 4 C.
- **Life cycle:** Infectious elementary bodies attached to the host mucosa and promoting its entry.. Cytoplasm phagosome.. producing reticulate bodies in inclusion.. released elementary bodies..
- **Chlamydia trachomatis..Serotypes C ,K** : Common cause of sexually transmitted disease (STD) Nonspecific urethritis.. mother to newborn babies..maternal fluid.. Atypical pneumonia..Eye infection..**Ophthalmia neonatorum**
- About half of all newborns with Chlamydial pneumonia develop inclusion conjunctivitis.. 1-2 weeks starts mild - severe eyes redness, swollen eyelids, inflammation & yellow thick discharge eyes.
- A & C serotypes of endemic *Ch. trachomatis* cause **Trachoma**.. conjunctival scarring, damage eyelids & Cornea.. blindness.

Chlamydia Life Cycle



Chlamydophila Pneumonia

- *C. pneumoniae*: droplets infection.. Infants/children often develops gradually.. several weeks mild respiratory symptoms, dry irritating prolonged cough.. nasal congestion.. with/without fever.. Few weeks.. No blood sepsis.
- *C. pneumoniae* infections in adults.. often asymptomatic, mild, May include sore throat, headache, fever, dry cough.
- Clusters of infection have been reported more common in Children than Adults.
- **Diagnosis & treatment:** Sputum, throat-nasal swab..
MaCoy Cell Culture, ELSA Specific antibodies, PCR and **Microimmunofluorescence MIF**.
- Treatment: Tetracyclines, Macrolides, levofloxacin, moxifloxacin .. No Vaccine

Chlamydophila Psittaci

- *C. psittaci* causes Zoonotic diseases.. Human infection followed contact with birds (parrots, pigeons, turkeys, and ducks).. A rare human disease called **psittacosis (ornithosis)**.
- Humans respiratory tract can be infected via inhalation bacteria shed from feathers, secretions, and droppings localized inflammation in Bronchi & lung tissues.
- Signs Symptoms: Starts mild..flu-like & ended with severe disease including fatal pneumonia, associated high fever, dry cough, headache.
- Diagnosis & Treatment similar to other Chlamydia.

Legionella pneumophila

- **Legionella** Gram negative , Pathogenic-Nonpathogenic spp. often found in natural aquatic bodies and wet soil. Facultative Anaerobes Growth in Cold/Hot (4- 80C) Water.. Transmitted, Inhalation via Air Condition, Wet Soil.. Cause outbreak of disease.
- Lung Mucosa..multiply intracellular within the macrophages.. High Fever .. Incub. period 2-10 days .. Nonproductive /Productive dry cough.. Shortness of breath, Chest pain, Muscle aches, Joint pain, Diarrhea, Renal Failure, higher mortality rate. Legionnaires' disease is not contagious
- **Risk factors** include heavy cigarette smoking, Old age underlying diseases such as **renal failure, cancer, diabetes, or** chronic obstructive pulmonary, suppressed immune systems, corticosteroid.
- **Diagnosis & treatment:** Special Culture Media, blood/urine specimen for detection Specific antibodies or Antigens by PCR, or EISA .. Macrolides (azithromycin), levofloxacin, moxifloxacin .. No Vaccine.

OPPORTUNISTIC MYCOSES

- Opportunistic mycoses are caused by globally distributed fungi that are either members of the human microbiota, such as a *Candida* species, or environmental yeasts and molds.
- They can produce disease ranging from superficial skin or mucous membrane infections to systemic involvement of multiple organs.
- Patients at risk include those with hematologic dyscrasias (eg, leukemia, neutropenia), patients with HIV/AIDS with CD4 counts less than 100 cells/ μ L, as well as those treated with immunosuppressive (eg, corticosteroid) or cytotoxic drugs

Cryptococcus neoformans

- *Cryptococcus neoformans* causes cryptococcosis.
- A widespread **encapsulated yeast** that inhabits soil around pigeon roosts
- Common infection of **AIDS, cancer** or **diabetes patients**
- Infection of **lungs** leads to cough, fever, and lung nodules
- **Dissemination to meninges** and brain can cause severe neurological disturbance and death.

Diagnosis

Microscopic

- India Ink for capsule stain (50-80% + CSF)

Culture

- Bird seed agar
- Routine blood culture

PCR

Aspergillosis: Diseases of the Genus *Aspergillus*

- Very common airborne soil fungus
- 600 species, 8 involved in human disease; *A. fumigatus* most commonly
- Serious opportunistic threat to **AIDS, leukemia, and transplant patients**
- Infection usually occurs in **lungs** – spores germinate in lungs and form **fungal balls**; can colonize **sinuses, ear canals, eyelids, and conjunctiva**
- **Bronchopulmonary allergy or Invasive aspergillosis in preformed cavities** can produce **necrotic pneumonia, and infection of brain, heart, and other organs.**
- Surgery , Amphotericin B and nystatin

Zygomycosis

- Zygomycota are extremely abundant saprophytic fungi found in soil, water, organic debris, and food.
- Genera most often involved are ***Rhizopus, Absidia, and Mucor.***
- Usually harmless air contaminants invade the membranes of the **nose, eyes, heart, and brain** of people (Rhinocerebral mucormycosis) with **diabetes** and malnutrition, with severe consequences.
- **main host defense is phagocytosis**

Diagnosis is made by direct smear and by isolation of molds from respiratory secretions or biopsy specimens.

Treatment:

Control Diabetes ,surgery &
amphotericin B

Prognosis: very poor

PNEUMOCYSTIS

- Pneumocystis jirovecii is the cause of a lethal pneumonia in immunocompromised persons, particularly those with AIDS.
- Definite diagnosis of pneumocystosis depends on finding organisms of typical morphology in appropriate specimens (Sputum, BAL)
- The organism has not been grown in culture
- TMP-SMX is treatment of choice

Endemic mycosis



- Endemic mycosis is caused by a thermally dimorphic fungus, and the infections are initiated in the lungs following inhalation of the respective conidia.
- Each of the four primary systemic mycoses—coccidioidomycosis, histoplasmosis, blastomycosis, and paracoccidioidomycosis—is geographically restricted to specific areas of endemicity.
- Most infections are asymptomatic or mild and resolve without treatment. However, a small but significant number of patients develop pulmonary disease.

Dimorphic Fungus: Histoplasmosis-1

- ***Histoplasma capsulatum***.. Dimorphic fungus with conidia and yeast forms at body temperature and hyphae & macroconidia in vitro culture.. Common in soil enriched with excreta of birds. Endemic in southern U.S.A, Australia.. Less other countries.
- The primary site of infection is usually pulmonary.. inhalation dust with microconidia.. Phagocytosed by macrophages, obligate intracellular parasites.. Causing slight inflammatory reaction.. Most cases of **histoplasmosis** are asymptomatic /subclinical, benign.. Flu-like syndrome.
- Few may develop chronic **progressive lung disease**.. Granuloma & fibrosis, chronic cutaneous or systemic disease involve any internal organ.. Fatal systemic disease.
- All infected persons become positive by histoplasmin skin test.

Histoplasma capsulatum in infected White Blood cells

H & E stained peripheral blood smear of a patient showing the yeast phase of *Histoplasma capsulatum* within white blood cells. MMRC-UTMB 1996



Coccidioidomycosis & Blastomycosis

- ***Coccidioides immitis* & *Blastomyces dermatitidis***.. soil inhabiting **Dimorphic Fungus**.. Endemic in south-western U.S.A., northern Mexico and various parts South America.
- Respiratory infection, resulting from the inhalation of microconidia, often resolves rapidly leaving the patient with a strong specific immunity to re-infection.
- Some individuals the disease may progress to a chronic **pulmonary** condition or a **systemic disease** involving the meninges, bones, joints, subcutaneous, cutaneous tissues.. Antigen Skin test positive.. Not significant in diagnosis.

Laboratory Diagnosis

- **Direct microscopy and culture** should be performed on all specimens (sputum, bronchial washings, CSF, pleural fluid tissue biopsies from various visceral organs).
- wet mounts in 10% KOH with india ink.. Ovoid-budding yeast cells (b) Gram-stain smear..
- Cultures on **Sabouraud dextrose agar** should be maintained for one month at 25C.... fungal growths & Wet Mount.. Identification ..produces hyphae-like conidio-phores & Spores.. Color of fungal growth
- **Serological tests are of limited value..** not significant
- Detection of Histoplasma antigen in blood & urine is significant

Paracoccidioidomycosis

- *Paracoccidioides brasiliensis* is the thermally dimorphic fungal agent of paracoccidioidomycosis (South American blastomycosis), which is confined to endemic regions of Central and South America.
- *P. brasiliensis* is inhaled, and initial lesions occur in the lung. After a period of dormancy that may last for decades, the pulmonary granulomas may become active, leading to chronic, progressive pulmonary disease or dissemination.

The End