

Doctor 2020

Collected by:
Laith Sami

1-Which one of the following isn't a criterion from Child Pugh scale? A-Albumin

C-Hemoglobin

D-PPT

B-Ascites

E-Bilirubin

Ans: C

2-Which of the following grade system is used to define severity of the esophageal reflux (GERD)?

A-Chicago grade

B-LA grade

C-Tokyo grade

D-Child Pugh grade

E-London grade

Ans: B

3-Which one of the following tests is specific for diagnosis of celiac disease?

A-IgA and IgG antigliadin antibodies

B-Anti reticulin antibodies

C-IgA and IgG tissue transglutaminase antibodies

D-IgA endomysial antibodies

E-Anti gluten antibodies

Ans: C

4-Which one of the following tests is specific for diagnosis of Hepatitis C?

A-Hepatitis C surface antigen

B-Hepatitis C core antigen

C- Total antibody to hepatitis C core antigen

D-Hepatitis C antibodies

Ans: D

5-Which of the following disease doesn't cause chronic diarrhea?

A-Giardiasis

B-Rota virus infection

C-HIV

D-Crohn's disease

E-celiac disease

Ans: B

6-Which one of the following drugs can't use for management of inflammatory bowel disease?

A- Azathioprine

B- Infliximab

C- Mercaptopurine

D- Non-Steroidal anti-inflammatory drugs

E- Methotrexate

Ans: D

7-Which one of the following approaches can't use for patient with Acute Esophageal varices?

A-IV Antibiotics

B-IV Fluid

C-IV PPI infusion

D-IV Hydrocortisone

E-Endoscopic rubber band ligation

Ans: D

8-Which of the following disease is common in Ulcerative Colitis patients?

- A- Primary Sclerosing Cholangitis
- **B-** Primary Biliary Cirrhosis
- C- Hemochromatosis
- **D-** Sarcoidosis
- E- All of the above

Ans: A

9-Which of the following is mostly to be found in patient with Budd Chari Syndrome?

A-High SSAG, Low protein

B-Low SSAG, Low protein

C-High SSAG, High protein

D-Low SAAG, High protein

E-Low SAAG, Normal protein

Ans: C

10-All of the following can cause esophageal dysphagia except?

A-Type 2 achalasia

B-Multiple sclerosis

C-Pseudo-achalasia

D-Esophageal varices

E-Esophageal cancer

Ans: B

11-Which of the following signs can't be found in a patient with Liver Cirrhosis?

- A- Spider angiomata
- B- Parotid gland atrophy
- C- Testicular atrophy
- **D- Splenomegaly**
- E- Clubbing and hypertrophic osteoarthropathy

Ans: B