



# Surgical approach to pathological thyroid gland


---

PROF. AYMAN MISMAR

# Introduction: Anatomy of the Thyroid Gland



**Anatomy of the Thyroid and Parathyroid Glands**





# Goals of Surgery:

- 1-to **remove** the primary tumour and its local extension.
- 2-to minimize treatment related **morbidity**.
- 3-to permit accurate **staging**.
- 4-facilitate postop. **Radioactive Iodine** ttt.
- 5-facilitate long term postop. **Surveillance**
- 6-minimize disease **recurrence** and mets.




□ Type of Surgery Vs approach.

ويعلمكم  
الكتاب والحكمة

١٩٦٢م ١٣٨٢

# Thyroidectomy types

## Types of Thyroidectomies








# Thyroidectomy types


- **Near-total thyroidectomy:** Almost same as total, but a little thyroid tissue around one parathyroid gland is preserved
- **Isthmusectomy:** Dividing the isthmus

# Conventional thyroidectomy











# Minimally Invasive thyroidectomy




# Miccoli: central neck access

: J Endoc Invest 1999



# Shimizu

Neck access : J Am Coll Surg 1999




**External Retraction  
(Kirschner)  
lateral Incision  
(SCM border)  
subclavicular Incision (5cm)**

Ikeda

axillary access: J Am Coll Surg

2000



**Incision 3 cm in axilla**

**Insufflation of CO<sub>2</sub> (4 mm Hg)**

**Flexible Endoscope**


**1 trocar near the main incision**



# Ohgami

*breast access* :


**Surg Laprosc 2000**



**Three incisions:**  
**1 presternal**  
**2 periareolar**  
**Insufflation of CO<sub>2</sub>**


# Gagner

*supraclavicular access* : *Thyroid 2001*





**Insufflation of CO<sub>2</sub> (8 mm Hg)**  
**Incision central (5 mm trocar)**  
**3 Trocars accessory:**  
**mid line**  
**mid border SCM**  
**sup border SCM**

# Delbridge 2006










# Terris


Robotic Face lift (retroauricular approach) : Laryngoscope 2011
















# **ENDOSCOPIC THYROIDECTOMY :**


## ***Different Techniques and Approaches***



- **Axillary and breast access should not be considered minimally invasive operations to approach thyroid**


**JF Henry Brit J Surg 2006**

# Wilhelm 2011 (transoral approach)













# Transoral Endoscopic Thyroidectomy: A Systematic Review of the Practice So Far

Christian Camenzuli, MD, Pierre Schembri Wismayer, MD, PhD, Jean Calleja Agius, MD, PhD

## ABSTRACT

**Background and Objectives:** Thyroid disease largely affects young females, but the incidence is also increasing among males. In an effort to avoid the scarring of the neck that is synonymous with conventional thyroidectomy, endoscopic techniques have been developed over the years. The transoral endoscopic approach is the latest of these innovations that promises a scarless surgical outcome. This review evaluates whether this technique is safe and feasible in live patients and outlines the outcomes in published literature so far.

**Database:** PubMed, Medline, BioMed Central, Cochrane Library, OVID and Web of Science were systematically searched by using a Medical Subject Heading (MeSH)-optimized search strategy. The selection of papers followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines after setting strict inclusion and exclusion criteria. Sixteen studies were included in the final analysis.

**Discussion:** This systematic review presents cases of 785 patients. Surgeons in 15 of the studies used a completely vestibular approach, whereas those in the remaining 2 used the floor of the mouth for primary access. Conversion to open surgery took place in 1.3%. In total, 4.3% of patients experienced transient laryngeal nerve palsy, whereas 0.1% had permanent recurrent incidences of the condition. Transient hypocalcemia occurred in 7.4% of cases, with no recorded permanent cases. Carbon dioxide embolism occurred in 0.6% of cases, and another 0.6%

had a deep-seated neck infection. The complication rates within the review were deemed acceptable and the overall technique feasible. A prospective randomized controlled trial was proposed to compare this technique with conventional thyroidectomy.

**Key Words:** Natural orifice endoscopic surgery, Oral endoscopy, Scarless, Thyroidectomy, Transoral.

## INTRODUCTION

Over the past centuries, procedures to surgically remove all or part of the thyroid gland from the neck have gone from infamy to fame. What Samuel D. Gross in the 19th century considered to be "horrid butchery," through the brave work of surgeons like Emil Theodor Kocher, has become one of the most common and safest of surgeries.<sup>1-3</sup> The gold-standard approach for thyroidectomy has been open or conventional surgery. Recently, there has been increased interest in applying the principles of minimally invasive surgery to thyroid surgery. This development was initially promoted by Miccoli and his colleagues<sup>4</sup> in 1999 and has continued to expand and improve throughout recent years. The aims of minimally invasive surgery include better cosmesis and earlier recovery without compromising the excellent results achieved with open surgery.<sup>5</sup> The approaches taken in thyroid surgery include mainly a transaxillary approach with later additions of areolar, anterior chest wall, and mixed approaches.<sup>6-10</sup> The extent of dissection and difficulty of these procedures despite robotic help has limited the uptake of these techniques.<sup>11,12</sup>

The transoral endoscopic technique, an adaptation of the concept of natural orifice transluminal endoscopic surgery (NOTES) to the neck, is a technique that promises to improve the aesthetic aspect by offering a scarless operation while retaining the advantages of minimally invasive surgery.<sup>13,14</sup> The pioneers of this technique were the group led by Witzel and his colleagues,<sup>15</sup> who presented their first paper on the subject in 2008. In their study on cadavers and live pigs, they managed to present a proof of concept that formed the basis for the extensive work that is being carried out by multiple groups around the world.

Department of Anatomy, Faculty of Medicine and Surgery, Biomedical Sciences, University of Malta, Msida, Malta. (all authors).

Disclosures: none reported.

Informed consent: Dr. Camenzuli declares that written informed consent was obtained from the patient/s for publication of this study/report and any accompanying images.

The authors thank Ms. Andee Agius and Ms. Roberta Sultana for proofreading the text and Endeavour Scholarship Malta for funding.

Address correspondence to: Christian Camenzuli, MD, Anatomy, Faculty of Medicine and Surgery, Biomedical Sciences Building, University of Malta, Msida Malta. Tel: +35679257480; E-mail: christian.camenzuli.04@um.edu.mt

DOI: 10.4293/JSLs.2018.00026

© 2018 by JSLs, Journal of the Society of Laparoendoscopic Surgeons. Published by the Society of Laparoendoscopic Surgeons, Inc.



# Till 2018

- 16 Reported series with a sum of 785 patients of which 713 (91%) were female and 68 (8%) were male.
- Anuwong et al (Thailand) 422 pts.
- Wilhelm et al (Germany) 96 pts.
- Fu J et al (China ) 81 pts.
- Yang J et al (China ) 46 pts.
- Other centers 130 pts.





- This systematic review presents cases of 785 patients. Surgeons in 15 of the studies used a completely vestibular approach, whereas those in the remaining 2 used the floor of the mouth for primary access. Conversion to open surgery took place in 1.3%. In total, **4.3%** of patients experienced **transient laryngeal nerve palsy**, whereas 0.1% had permanent recurrent incidences of the condition. Transient hypocalcemia occurred in 7.4% of cases, with no recorded permanent cases. Carbon dioxide embolism occurred in 0.6% of cases, and another 0.6% had a deep-seated neck infection.



الحكمة


# *Miccoli (MIVAT) 1998*



Patient in supine  
position  
Neck not extended  
Skin covered by  
drape



# MIVAT: Instrumentation





**30° 5mm endoscope**  
**Spatulas (2mm)**  
**Spatula-aspirator**  
**Scissors (2mm)**  
**Forceps (2mm)**  
**Retractors**  
**Clip applier**  
**Harmonic Scalpel**



# MIVAT Incision
















# ***INCLUSION CRITERIA***

- Nodule less than 3.5 cm**
  
- Thyroid volume less than 20 ml  
(ultrasound measured)**
  
- No lymphnodes**
  
- No thyroiditis**





# COMPLICATIONS

- **18 transient laryngeal nerve palsy (1.8 %)**
- **9 perman. laryngeal nerve palsy (0.9 %)**
- **18 transient hypoparathyroidism (2.2 %)**
- **3 permanent hypoparathyroidism (0.6%)**

**same rate in multicentric studies  
comparable to standard surgery**

## American Thyroid Association Statement on Remote-Access Thyroid Surgery (2016)



- Remote-access thyroidectomy has a role in a small group of patients who fit strict selection criteria. These approaches require an additional level of expertise, and therefore should be done by surgeons performing a high volume of thyroid and robotic surgery.



# Pros

- Cosmetic outcome.

- Tissue injury.

# Cons




- ❑ Visualization and field.
- ❑ Tactile sensation.
- ❑ Transient complications.
- ❑ Remote dissection.
- ❑ Oncological outcome.
- ❑ Cost effectiveness.



# Nerve monitor







Thank You