



Psychiatry Mini-OSCE

Doctor 2020 Mini-OSCE

Special thanks to

1st rotation:

Ammar Ali

Alaa Mo'taz

Noor muneer

Noor sufian

2nd rotation:

Insaf Alammouri

Ahmad Riyad

Fawzi Qadoomi

Bayan Shiekh Ali

1st rotation

Case 1

Woman came to ER with panic attack, calmed down with breathing technique vital signs was normal (HR normal). History shows she had a job interview?. She has always been shy in public, and afraid to say silly things. She doesn't expect another panic attack.

1. What is the diagnosis?

Social phobia.

2. Does she meet criteria for panic disorder?

No.

3. She was given SSRI (escitalopram), which drug should be given with it?

Benzodiazepines.

4. She came the next day with another panic attack, why?

Due to side effect (anxiety) of SSRI and need time to exert the effect.

5. If a patient had fear of public speaking, what medication do you give?

B- blockers.

Case 2

Patient brought by concerned family, very talkative over enthusiastic, exchanged numbers with attractive lady, buys a lot of things, stays up at night doesn't sleep, talks on phone... Had a depressive episode 5 years ago.

1. Diagnosis?

Bipolar 2.

2. Medication to give?

Lithium.

3. Which medication should be avoided?

Antidepressants.

4. Is there a genetic relationship with this disease?

Yes.

5. If patient with same current episode came the following year? With psychosis, what will his diagnosis be?

No (it is not mania /no functional impairments).

Case 3

Patient found walking on the streets talking to himself doing wierd inappropriate signs, when police confronted him he ran away yelling "ما حخلي العصابات يمسوني (اشي زي هيك)" he had poor higene, long hair... Etc

1. Name the 3 phases of this condition?

Prodromal / active / residual.

2. Name the 3 categories of symptoms?

Positive, negative, cognitive.

3. Why is it more common in lower Socioeconomic groups?

Downward drift phenomenon / many patients face barriers to higher education, regular employment and other resources, so they tend to drift downward Socioeconomically.

4. Which have better prognosis males or females?

Females.

5. Compare old and new medications side effect profile?

Typical antipsychotics - extra pyrimdial symptoms / atypical antipsychotics metabolic.

Case 4

Patient brought by his friends, he went to a party with them, took a substance to help him feel energized and stay up all night. After a while he started hearing voices and seeing things. Became paranoid of ghosts chasing him.

1. What class of drugs does the substance belong to?

Stimulants.

2. Are pupils dilated or constricted in patients taking this substance?

Dilated.

3. What neurotransmitters is increased?

Dopamine.

4. What medication is given in intoxiaction?

Benzodiazepines.

5. Is sudden stopping of drug life threatening?

No.

6. In withdrawl is patient sedated or aroused?

Sedated.

7. Is there antidote?

No.

Case 5

Patient revisiting clinic asked you the following

1. Which 2 organs are affected by lithium?

Kidney / thyroid.

2. Lethal serum level of lithium?

2 mEq/L.

3. Therapeutic serum level of valproic acid?

80 -120 mg/L.

4. 2 classes of medication that increase serum level of lithium?

Thiazide diuretics:

- NSAIDs.
- ACE inhibitors.
- Metronidazole.
- Tetracycline.

5. Life threatening side effect of Lamotrigine?

Stevens-Johnson syndrome.

6. Teratogenic effect of anticonvulsants in pregnancy?

Neural tube defects.

2nd rotation

Case 1

A 70 year old hospitalized patient was checked for depression. He has HTN, DM, 2 CVAs, HF=25%, was admitted for suspected pneumonia and on IV antibiotics, annoyed from the dog sounds in the room although dogs are not allowed in the hospital, and was found trying to remove his foley's cath.

1. What is his diagnosis?

Delirium.

2. What is the appropriate management?

Treat underlying cause, antipsychotics, family supervision.

3. Should we give patients with delirium benzo? And why?

No, it causes paradoxical disinhibition or over sedation, and patient has depression so you can't give benzo.

4. What's the difference between dementia and delirium (mention 2)?

Dementia: Chronic progressive decline in cognitive functions, preserved level of consciousness, usually irreversible (except if caused by vitamin B12 deficiency, hypothyroidism), normal EEG.

Delirium: Acute, waxing and waning level of consciousness, reversible, diffuse background slowed pattern EEG.

5. mention 2 risk factors of delirium?

Old age, comorbid diseases, prior history of delirium.

6. When does it become worse?

At night (sundowning).

Case 2

A 6 year old boy is brought by his mother, concerned he hasn't said "mama, dada" yet, quiet, prefers to play alone, doesn't get along with his classmates, annoyed when his toys are not in the order he's used to. In the clinic, you call him and he doesn't answer and acts as if you're not there.

1. What is the diagnosis?

Autism spectrum disorder.

2. What is the treatment?

a-2 agonists, early intervention, remedial education, behavioral therapy, psychoeducation, but the question asked for the "curative treatment" so make sure to mention that there is no curative treatment for autism.

3. What are the changes that may be seen when he becomes an adult?

Inrelectual function and language impairment.

4. More in Male or females?

Males.

5. Aside from language testing, what other medical test should you perform?

Formal Neuropsychological testing, Auditory testing.

6. What is the most common single genetic mutation associated with the disease?

Fragile x syndrome.

Case 3

A 23 year old female is concerned about her nose shape and size and is afraid of what people might say about it, and says it looks like a carrot on my face. She thinks that surgery will be the only way to fix it, examination showed a normal size and shape for woman, and her family says they see it normal.

1. What is the diagnosis?

Body dysmorphic disorder.

2. More common in Male or female?

Females.

3. Associated disorders?

MDD, OCD, social anxiety.

4. Satisfaction with plastic surgeries?

No.

5. High risk of suicide?

Yes.

6. Management?

SSRI, CBT.

Case 4

A 40 year old pt was admitted because he has thoughts of killing his brother everytime he sees him, he says that he avoids seeing him so his thoughts don't come back.

1. What is the diagnosis?

Ocd, intrusive taboo thoughts.

2. Other subtypes?

Contamination, doubt or harm, symmetry.

3. Is it better to tell the family about his thoughts?

Yes, family support can help the patient through his treatment plan and to protect the patient and others.

4. Pharmacological treatment of choice?

SSRIs.

5. Type of psychotherapy?

CBT (Exposure and Response Prevention).

6. Does it have a strong genetic factor?

Yes.

Case 5

1. Cluster C types?

Avoidant, dependance, ocpd.

2. Schizoid vs avoidant?

Schizoid: cluster a, psychotic symptoms, prefer to be alone, while avoidant is cluster C, anxiety symptoms, prefer to be with people but are too shy.

3. Do patients have insight ?

Patients often lack insight.

4. Type of psychotherapy?

cbt.

5. Can they have criteria of more than one personality disorders?

Yes, multiple personality disorders can coexist.

6. Antisocial pts have as children?

Conduct disorder.